


Table 5.14
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2005-2009

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.39	0.02		
High Five States				
Mississippi	3.88 ^b	0.22	(01)	62.23 ^c
Louisiana	3.35 ^b	0.17	(02)	40.17 ^c
Arkansas	3.32 ^b	0.21	(03)	39.10 ^c
Alabama	3.04 ^b	0.15	(04)	27.32 ^c
D.C.	2.94	0.43	(05)	23.07 ^c
Low Five States				
Minnesota	1.49 ^b	0.10	(47)	-37.51 ^c
Rhode Island	1.45 ^b	0.21	(48)	-39.20 ^c
Massachusetts	1.39 ^b	0.09	(49)	-41.90 ^c
Utah	1.30 ^b	0.16	(50)	-45.74 ^c
Vermont	1.14 ^b	0.25	(51)	-52.17 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.04 ^b	0.15	(04)	27.32 ^c	Montana	1.94	0.28	(36)	-18.64 ^c
Alaska	1.99	0.39	(34)	-16.66 ^c	Nebraska	1.59 ^b	0.18	(45)	-33.64 ^c
Arizona	1.90 ^b	0.11	(41)	-20.44 ^c	Nevada	2.69	0.21	(12)	12.61
Arkansas	3.32 ^b	0.21	(03)	39.10 ^c	New Hampshire	2.09	0.24	(32)	-12.71
California	2.35	0.05	(23)	-1.68	New Jersey	2.44	0.10	(20)	2.12
Colorado	1.74 ^b	0.12	(42)	-27.26 ^c	New Mexico	2.65	0.23	(15)	10.97
Connecticut	1.64 ^b	0.13	(44)	-31.34 ^c	New York	2.38	0.07	(22)	-0.34
Delaware	2.18	0.30	(27)	-8.67	North Carolina	2.35	0.10	(24)	-1.88
D.C.	2.94	0.43	(05)	23.07 ^c	North Dakota	1.54	0.30	(46)	-35.65 ^c
Florida	2.68 ^b	0.07	(13)	12.17	Ohio	2.59	0.09	(17)	8.37
Georgia	2.68	0.11	(14)	12.16	Oklahoma	2.91	0.18	(07)	21.63 ^c
Hawaii	1.93	0.24	(37)	-19.29 ^c	Oregon	1.92	0.14	(38)	-19.78 ^c
Idaho	2.14	0.24	(30)	-10.46	Pennsylvania	2.25	0.08	(26)	-5.72
Illinois	2.63	0.09	(16)	9.83	Rhode Island	1.45 ^b	0.21	(48)	-39.20 ^c
Indiana	2.38	0.12	(21)	-0.31	South Carolina	2.85	0.15	(10)	19.03 ^c
Iowa	2.11	0.16	(31)	-11.87	South Dakota	1.91	0.30	(39)	-20.18 ^c
Kansas	2.18	0.18	(28)	-8.81	Tennessee	2.79	0.13	(11)	16.79 ^c
Kentucky	2.91	0.16	(06)	21.92 ^c	Texas	2.91 ^b	0.07	(08)	21.59 ^c
Louisiana	3.35 ^b	0.17	(02)	40.17 ^c	Utah	1.30 ^b	0.16	(50)	-45.74 ^c
Maine	1.90	0.22	(40)	-20.34 ^c	Vermont	1.14 ^b	0.25	(51)	-52.17 ^c
Maryland	2.28	0.12	(25)	-4.77	Virginia	2.18	0.10	(29)	-8.89
Massachusetts	1.39 ^b	0.09	(49)	-41.90 ^c	Washington	1.96 ^b	0.11	(35)	-18.16 ^c
Michigan	2.00 ^b	0.09	(33)	-16.17 ^c	West Virginia	2.90	0.24	(09)	21.51 ^c
Minnesota	1.49 ^b	0.10	(47)	-37.51 ^c	Wisconsin	1.70 ^b	0.11	(43)	-28.98 ^c
Mississippi	3.88 ^b	0.22	(01)	62.23 ^c	Wyoming	2.53	0.45	(19)	6.01
Missouri	2.57	0.13	(18)	7.71					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.